

GASTROPANEL® EXAMINATION

fP-GastPan (9172)

The study examines upper abdominal symptoms, gastric mucosal damage and related risks, as well as possible *Helicobacter pylori* infection and an acid-free stomach.

How to prepare for the examination

A blood test is taken in the laboratory, before which the patient **must not eat, drink or smoke for 4–10 hours**.

Before sampling, certain medicine should be avoided. **Any discontinuation of medication should always be discussed with a doctor.**

Medicine that cannot be taken for 24 hours before the examination

- Hydrochloric acid secretion inhibitors: Cytotec, Famotidin Stada, Pepcid, Ranisan, Ranitidin Mylan, Ranixal, Zantac, Inside Brus, Ranitidin-Ratiopharm
- Agents impacting the intestinal motility: Constella, Gaspid, MCP-Ratiopharm, Metoclopramide Accord, Metoclopramide Orion, Motilium, Motilium Takeda, Motilium Tropfen, Primperan, Resolor, Mestinon, Ubretid Takeda Pharma
- Antacids: Balancid Novum, Magnesiamaito, Rennie, Natriumbikarbonat Recip
- Cytoprotective agents: Antepsin, Galieve, Gaviscon

Medicine that cannot be taken for two (2) weeks before the examination

- Courses of antibiotics
- Proton pump inhibitors: Acidal, Esomeprazol Actavis, Esomeprazol Krka, Esomeprazol Orion, Esomeprazol Ratiopharm, Esomeprazol Sandoz, Gasterix, Lanrec, Lansoprazol Krka, Lansoprazol Mylan, Lansoprazole, Lantil, Lanvone, Losec, Nexium, Omeprazol Actavis, Omeprazol Ratiopharm, Omeprazol Sandoz, Omestad, Pantopratsoli Sun, Pantoprazol Actavis, Pantoprazol Krka, Pantoprazol Mylan, Pantoprazol Orion, Pantoprazole Ratiopharm, Pariet, Rabeprazol Krka, Somac, Somac Control, Zolt.

Medicine that cannot be taken for four (4) weeks before the examination

- *Helicobacter pylori* eradication therapy: Helipak A, Helipak K

If, for any reason, the use of the above-mentioned medicines affecting gastric secretion cannot be interrupted, this should be mentioned in the referral.

The sample must be accompanied by information on *Helicobacter pylori* eradication treatment, heartburn symptoms and the use of anti-acidity and analgesic medicines. Notify the laboratory, if the instructions have not been followed.